

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Modeling epidemic spreading in star-like networks

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/127882> since

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Modeling epidemic spreading in star-like networks

Luca Ferreri, Paolo Bajardi, Mario Giacobini

GECO - Group of Computational Epidemiology
Department of Veterinary Sciences


CSU - Complex Systems Unit
Molecular Biotechnology Center

ARC²S - Applied Research on Computational Complex Systems Group
Department of Computer Science University of Torino


Milano, 5 aprile 2013

Tick-Borne Encephalitis


- ▶ endemic in Eurasia from Europe, through Russia To China and Japan
- ▶ the virus causes potentially fatal neurological infection
- ▶ in last years emergence of the virus in new area and increase of morbidity
- ▶ maintained in nature by complex cycle involving Ixodid ticks (*I. ricinus* and *I. persulcatus*) and wild vertebrate hosts



Systemic Transmission




time t




time $t + 1$


Non-Systemic Transmission



Non-Systemic Transmission



Non-Systemic Transmission




Our research question

how do the **non-systemic transmission** together with the **different aggregation patterns** influence the pathogen spreading?


Spreading Model

- ▶ at time t a fraction, $\pi(t)$, of passengers (ticks) are infectious




Spreading Model

- ▶ at time t a fraction, $\pi(t)$, of passengers (ticks) are infectious
- ▶ $\mathbb{P}(k)$ probability that a bus (mouse) transports k passengers (ticks) of them




Spreading Model

- ▶ at time t a fraction, $\pi(t)$, of passengers (ticks) are infectious
- ▶ $\mathbb{P}(k)$ probability that a bus (mouse) transports k passengers (ticks) of them



Spreading Model

- ▶ at time t a fraction, $\pi(t)$, of passengers (ticks) are infectious
- ▶ $\mathbb{P}(k)$ probability that a bus (mouse) transports k passengers (ticks) of them
- ▶ β transmission probability for infectious path
- ▶ μ recovery probability





Spreading Model

- ▶ at time t a fraction, $\pi(t)$, of passengers (ticks) are infectious
- ▶ $\mathbb{P}(k)$ probability that a bus (mouse) transports k passengers (ticks) of them
- ▶ β transmission probability for infectious path
- ▶ μ recovery probability

$$\Rightarrow \pi(t+1) = f(\pi(t))$$

time $t + 1$



Analytical Framework

the probability that a **susceptible passenger**, having h travel mates, gets the **infection** is

$$1 - (1 - \beta)^h$$

Analytical Framework

the probability that a **susceptible passenger**, having h travel mates, gets the **infection** is

$$1 - (1 - \beta)^h$$

Let $\pi(t)$ be the **prevalence of infection** among passengers at time t , the probability for a susceptible passenger on a bus transporting k individuals including himself to be **infectious** at time $t + 1$ is

$$1 - (1 - \beta)^{(k-1) \cdot \pi(t)}$$

Math

Recalling that $\mathbb{P}(k)$ is the probability for a bus to have k passengers, the probability for a passenger to be on a k -bus is

$$\begin{aligned}\frac{\text{\#passengers on a } k\text{-bus}}{\text{\#passengers}} &= \frac{k \cdot \text{\#}k\text{-bus}}{\text{\#passengers}} = \\ &= k \cdot \frac{\text{\#}k\text{-bus}}{\text{\#bus}} \cdot \frac{\text{\#bus}}{\text{\#passengers}} = k \cdot \mathbb{P}(k) \cdot \frac{1}{\langle k \rangle}\end{aligned}$$

thus, the probability for a **susceptible** passenger at time t to be **infectious** at time $t + 1$ is

$$\sum_{k=1}^{\infty} \left[1 - (1 - \beta)^{(k-1) \cdot \pi(t)} \right] \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k)$$

and therefore the prevalence among passenger at time $t + 1$ is

$$\begin{aligned} \pi(t+1) &= f(\pi(t)) = \\ &= (1 - \mu) \cdot \pi(t) + [1 - \pi(t)] \cdot \left\{ 1 - \sum_{k=1}^{\infty} (1 - \beta)^{(k-1) \cdot \pi(t)} \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k) \right\} \end{aligned}$$

Equilibria

imposing the stationary condition $\pi(t+1) = \pi(t) = x$ we can derive the equilibria as solutions of the following equation

$$x = f(x) = (1 - \mu) \cdot x + [1 - x] \cdot \left\{ 1 - \sum_{k=1}^{\infty} (1 - \beta)^{(k-1) \cdot x} \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k) \right\}.$$

Equilibria

imposing the stationary condition $\pi(t+1) = \pi(t) = x$ we can derive the equilibria as solutions of the following equation

$$x = f(x) = (1 - \mu) \cdot x + [1 - x] \cdot \left\{ 1 - \sum_{k=1}^{\infty} (1 - \beta)^{(k-1) \cdot x} \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k) \right\}.$$

Now:

- ▶ $x = 0$ is a solution,
- ▶ $f(1) = 1 - \mu \leq 1$,
- ▶ $f''(x) < 0$.

Equilibria


imposing the stationary condition $\pi(t+1) = \pi(t) = x$ we can derive the equilibria as solutions of the following equation

$$x = f(x) = (1 - \mu) \cdot x + [1 - x] \cdot \left\{ 1 - \sum_{k=1}^{\infty} (1 - \beta)^{(k-1) \cdot x} \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k) \right\}.$$

Now:

- ▶ $x = 0$ is a solution,
- ▶ $f(1) = 1 - \mu \leq 1$,
- ▶ $f''(x) < 0$.

assuming $\langle k \rangle$ and $\langle k^2 \rangle$ finite



Equilibria


imposing the stationary condition $\pi(t+1) = \pi(t) = x$ we can derive the equilibria as solutions of the following equation

$$x = f(x) = (1 - \mu) \cdot x + [1 - x] \cdot \left\{ 1 - \sum_{k=1}^{\infty} (1 - \beta)^{(k-1) \cdot x} \cdot \frac{k}{\langle k \rangle} \cdot \mathbb{P}(k) \right\}.$$

Now:

- ▶ $x = 0$ is a solution,
- ▶ $f(1) = 1 - \mu \leq 1$,
- ▶ $f''(x) < 0$.

assuming $\langle k \rangle$ and $\langle k^2 \rangle$ finite



therefore conditions to have one, and only one, solution $\hat{x} \in (0, 1)$ is that $f'(0) > 1$ or

$$-\frac{\ln(1 - \beta)}{\mu} > \frac{\langle k \rangle}{\langle k^2 \rangle - \langle k \rangle}.$$

Stability

recalling that $f''(x) < 0$ and that

- ▶ $f'(1) = -\mu + \sum_k (1 - \beta)^{k-1} \frac{k}{\langle k \rangle} \mathbb{P}(k) > -\mu > -1$
- ▶ $f'(\hat{x}) < 1$

hence \hat{x} is **asymptotically stable** when it exists. Therefore:

- ▶ disease-free equilibrium is asymptotically stable when \hat{x} does not exist.
- ▶ disease-free is unstable when \hat{x} exists. Furthermore, when \hat{x} exists it is also asymptotically stable.

Conclusion and Discussions

- ▶ co-feeding transmission
- ▶ spreading on star-like networks
- ▶ spreading dynamic on dynamic bipartite networks
- ▶ analytical result **confirmed** by simulations

Acknowledgements

Source of inspirations for this work come from a scientific project hold with the Fondazione Edmund Mach, San Michele all'Adige (TN). In particular, we are grateful to Anna Paola Rizzoli, Roberto Rosà e Luca Bolzoni.